

MELANIE PRINCE, MD

PLASTIC SURGERY

Nicotine Policy

Smoking patients, patients who use other nicotine products, and patients exposed to second-hand smoke, experience a decreased healing capacity after surgery. The complications these patients are at increased risk for after surgery include:

- Loss of skin, nipples or tummy skin after a breast lift, breast reduction, or tummy tuck surgery
- Infections
- Death of fat cells (fat necrosis), causing hard lumps
- Delayed wound healing
- Thick, wide scars
- Blood clots, which can be fatal
- Increased pain
- Permanent small vessel damage adding risk even if you quit
- Loss of breast implants
- Life-threatening complications like stroke, heart attack, blood clots, and pneumonia.
- Complications from or under anesthesia

We require that all patients completely avoid nicotine, any nicotine-containing products, and second-hand smoke a minimum of 6 weeks before surgery and up to 6 weeks after surgery. No nicotine or nicotine-containing products can be in your system during this time. These products include:

- Cigarettes
- Cigars, Little Cigars, Cigarillos
- Dissolvable Nicotine Products: Lozenges, mints, strips, or sticks
- Electronic Cigarettes
- Smokeless Tobacco Products: Chewing tobacco, moist snuff
- Waterpipes
- Nicotine Patches
- Nicotine Gum
- Marijuana rolled in cigar paper
- Second-Hand Smoke from any of the above

Initial

I understand all patients will be nicotine tested the day of surgery. If the test is positive, my surgery will be cancelled, all money will be forfeited, and the surgery will not be rescheduled.

Signature: _____

Date: _____

Printed Name: _____